

A relative Yoneda Lemma (manuscript)

M. Künzer

February 8, 2008

Abstract

We construct set-valued right Kan extensions via a relative Yoneda Lemma.

A remark of the referee

As the referee pointed out, (2.1) ‘can essentially be found in much greater generality’ in

[Ke 82] G. M. KELLY, *Basic concepts of enriched category theory*, LMS Lecture Notes 64, Cambridge University Press, 1982.

He continued to explain that to this end, one reformulates the formula [Ke 82, 4.6 (ii), p. 113], given in terms of weighted limits, by means of [Ke 82, 3.10, p. 99] and [Ke 82, 2.2, p. 58].

Therefore, we withdraw this note as a preprint. Since (2.1, 3.1) might be of some use for the working mathematician, we leave it accessible as a manuscript. In using (2.1), the reader should refer to [Ke 82], in using (3.1), he should refer to [K 58].

0 Introduction

0.1 A relative Yoneda Lemma

The category of set-valued presheaves on a category C shall be denoted by $C^\wedge$. The Yoneda embedding, sending x to ${}_C(-, x)$, shall be denoted by $C \xrightarrow{y_C} C^\wedge$. The presheaf category construction being contravariantly functorial, we obtain the

Proposition (2.1, the relative Yoneda Lemma). *Given a functor $C \xrightarrow{f} D$, we have*

$$f^\wedge \dashv y_D^\wedge \circ f^{\wedge\wedge} \circ y_{C^\wedge}.$$

There is a set theoretical caveat. In particular, this formula is correct only after some additional comments, see (2.1).

The right hand side is also known as the set-valued right Kan extension functor along $C^o \xrightarrow{f^o} D^o$. There are several formulas for right Kan extensions in the literature, for instance using ends [ML 71, X.4], all of them necessarily yielding the same result up to natural equivalence, by uniqueness of the adjoint. In particular, (2.1) is merely still another such a formula.

Letting $f = 1_C$, we recover the (absolute) Yoneda Lemma, thus giving a solution to the exercise [ML 71, X.7, ex. 2]. Concerning its origin, MAC LANE recalled the following incident, taking place in around 1954/55 [ML 98].

MAC LANE, then visiting Paris, was anxious to learn from YONEDA, and commenced an interview with YONEDA in a café at the Gare du Nord. The interview was continued on YONEDA's train until its departure. In its course, MAC LANE learned about the lemma and subsequently baptized it.

0.2 A relative co-Yoneda Lemma

Suppose given a functor $C \xrightarrow{k} Z$. KAN constructed in [K 58, Th. 14.1] the left adjoint to the functor from Z to $C^\wedge$ that sends $z \in Z$ to $\mathcal{Z}(k(-), z) \in C^\wedge$. Now given a functor $C \xrightarrow{f} D$, we can specialize to $Z = D^\wedge$ and to a functor k that sends $c \in C$ to $\mathcal{D}(-, fc) \in D^\wedge$, thus obtaining a left adjoint to $C^\wedge \xleftarrow{f^\wedge} D^\wedge$. (KAN's notation is as follows. Identify $\mathcal{Z} = Z$, $\mathcal{V} = C$, $\mathfrak{M}^\mathcal{V} = C^\wedge$. The functor $H^\mathcal{V}(\mathcal{Z}_\mathcal{V}, \mathcal{Z})$ maps from $\mathcal{Z}_\mathcal{V}, \mathcal{Z}$ to $\mathfrak{M}^\mathcal{V}$, i.e. from (the category of covariant functors from $\mathcal{V}$ to $\mathcal{Z}$) $\times$ $\mathcal{Z}$ to the category of presheaves on $\mathcal{V}$. Given $\mathcal{V} \xrightarrow{k} \mathcal{Z}$ and an object $z \in \mathcal{Z}$, the pair (k, z) is mapped to $\mathcal{Z}(-, z) \circ k$.) We shall rephrase this special case of KAN's formula as follows, in order to be able to compare the left and the right adjoint of $f^\wedge$.

Let $C^\vee := (C^o)^\wedge$. The co-Yoneda embedding, sending x to $\mathcal{C}(x, -)$, shall be denoted by $C \xrightarrow{y'_C} C^\vee$. There is a tensor product $C^\wedge \times C^\vee \longrightarrow (\text{Set})$, sending $v \times v'$ to $v \otimes_C v'$. The according univalent functor that sends v to $v \otimes_C -$ shall be denoted by $C^\wedge \xrightarrow{z_C} C^{\vee\vee}$.

Proposition ([K 58], cf. 3.1, the relative co-Yoneda Lemma). *Given a functor $C \xrightarrow{f} D$, we have*

$$y'_D{}^\vee \circ f^{\vee\vee} \circ z_C \dashv f^\wedge.$$

Again, there is a set-theoretical caveat.

0.3 Acknowledgements

I would like to thank F. BORCEUX for pointing out set theoretical difficulties. I would like to thank R. RENTSCHLER for kind hospitality. The author received support from the EU TMR-network 'Algebraic Lie Representations', grant no. ERB FMRX-CT97-0100.

0.4 Notation

- (i) We write composition of morphisms on the right, $\xrightarrow{a} \xrightarrow{b} = \xrightarrow{ab}$. However, we write composition of functors on the left, $\xrightarrow{f} \xrightarrow{g} = \xrightarrow{g \circ f}$.
- (ii) Given a functor $C \xrightarrow{f} D$, the opposite functor between the opposite categories is denoted by $C^o \xrightarrow{f^o} D^o$.
- (iii) Given functors $C \xrightleftharpoons[f']{f} D \xrightleftharpoons[g']{g} E$ and natural transformations $f \xrightarrow{\alpha} f'$ and $g \xrightarrow{\beta} g'$, we denote by $g \circ f \xrightarrow{\beta \circ f} g' \circ f$ the natural transformation defined by $(\beta \circ f)c = \beta(fc)$ for $c \in C$ and by $g \circ f \xrightarrow{g \circ \alpha} g \circ f'$ the natural transformation defined by $(g \circ \alpha)c = g(\alpha c)$ for $c \in C$. More generally,

$$(g \circ f \xrightarrow{\beta \circ \alpha} g' \circ f') := (g \circ f \xrightarrow{\beta \circ f} g' \circ f \xrightarrow{g' \circ \alpha} g' \circ f') = (g \circ f \xrightarrow{g \circ \alpha} g \circ f' \xrightarrow{\beta \circ f'} g' \circ f').$$

1 Universes

Since we will iterate the construction ‘forming the presheaf category over a category’ once, we shall work in the setting of universes, which enables us to do such ‘large’ constructions when keeping track of the universe needed. Therefore, we start with a preliminary section to recall this well-known technique from [SGA 4 I, App.] and to fix some notation.

Definition 1.1 (N. BOURBAKI, [SGA 4 I, App., 1, Déf. 1])

A universe is a set $\mathfrak{U}$ that satisfies the conditions (U1-4).

(U1) $x \in y \in \mathfrak{U}$ implies $x \in \mathfrak{U}$.

(U2) $x, y \in \mathfrak{U}$ implies $\{x, y\} \in \mathfrak{U}$.

(U3) $x \in \mathfrak{U}$ implies $\mathfrak{P}(x) \in \mathfrak{U}$.

(U4) Given $I \in \mathfrak{U}$ and a map $I \longrightarrow \mathfrak{U}$, $i \longmapsto x_i$, the union $\bigcup_{i \in I} x_i$ is in $\mathfrak{U}$.

Here $\mathfrak{P}(x)$ denotes the power set of x .

Remark 1.2

- (i) $x \in \mathfrak{U}$ implies $\{x\} = \{x, x\} \in \mathfrak{U}$.
- (ii) $x \subseteq y \in \mathfrak{U}$ implies $x \in \mathfrak{U}$. In particular, if y surjects onto some set z , then z is in bijection to an element of $\mathfrak{U}$.
- (iii) Let $(x, y) := \{x, \{x, y\}\}$. The element x is the unique element of

$$\{a \in (x, y) \mid \text{for all } b \in (x, y) \setminus \{a\} \text{ we have } a \in b\},$$

since $\{x, y\} \in x$ would contradict VON NEUMANN’s axiom, asserting that any nonempty set S contains an element that has an empty intersection with S . Now $\{x, y\}$ is the unique element of $(x, y) \setminus \{x\}$, and y is the unique element of $\{x, y\} \setminus \{x\}$.

- (iv) $X \in \mathfrak{U}$ and $Y \in \mathfrak{U}$ implies $X \times Y := \bigcup_{x \in X} \bigcup_{y \in Y} \{(x, y)\} \in \mathfrak{U}$.
- (v) Given $I \in \mathfrak{U}$ and a map $I \longrightarrow \mathfrak{U}$, $i \longmapsto x_i$, the disjoint union $\coprod_{i \in I} x_i := \bigcup_{i \in I} X_i \times \{i\}$ is in $\mathfrak{U}$.
- (vi) Given $I \in \mathfrak{U}$ and a map $I \longrightarrow \mathfrak{U}$, $i \longmapsto x_i$, the product $\prod_{i \in I} x_i \subseteq \mathfrak{P}(\coprod_{i \in I} x_i)$ is in $\mathfrak{U}$.

Assume given universes $\mathfrak{U}$, $\mathfrak{V}$ and $\mathfrak{W}$ such that

$$\emptyset \neq \mathfrak{U} \in \mathfrak{V} \in \mathfrak{W}.$$

This may be achieved by means of BOURBAKI's axiom (A.6) in [SGA 4 I, App., 4], which says that any set is element of some universe. Note that $\mathfrak{U} \subseteq \mathfrak{V} \subseteq \mathfrak{W}$.

For set theoretical purposes the following definition is convenient.

Definition 1.3 (cf. [M 65, I.2]) *A small category C is a tuple $(M, K \subseteq M \times M \times M)$, M being a set, subject to the conditions (i-iii). We write $M =: \text{Mor } C$, the set of morphisms of C . K is the composition law of C .*

- (i) *For every $(a, b) \in M \times M$ there exists at most one $c \in M$ such that $(a, b, c) \in K$. In this case, we write $c = ab$ and say that '(the composite) ab exists'.*

Let

$$\begin{aligned} \text{Ob } C := \{ a \in M \mid & \text{ for all } b \text{ for which } ba \text{ exists, we have } ba = b, \text{ and} \\ & \text{ for all } c \text{ for which } ac \text{ exists, we have } ac = c. \} \subseteq \text{Mor } C \end{aligned}$$

be the set of objects of C . If $a \in \text{Ob } C$, we also write $a =: 1_a$.

- (ii) *For every $a \in M$ there exists a unique source element $s(a) \in \text{Ob } C$ such that $s(a)a$ exists, and a unique target element $t(a) \in \text{Ob } C$ such that $at(a)$ exists.*
- (iii) *For $a, b \in M$ such that the composite ab exists, the composites $t(a)b$ and $as(b)$ exist.*
- (iv) *Given $a, b, c \in M$ such that ab and bc exist. Then $(ab)c$ and $a(bc)$ exist and equal each other.*

The set of morphisms with start x and target y shall be denoted by ${}_x C_y$.

Suppose given $a, b \in M$. Note that if ab exists, we have $s(ab) = s(a)$ and $t(ab) = t(b)$. Note that $s(s(a)) = s(a)$, $t(t(a)) = t(a)$. Note that ab exists iff $t(a) = s(b)$.

Definition 1.4 *A set X is said to be $\mathfrak{U}$ -small if there exists a bijection from X to an element of $\mathfrak{U}$. The category of $\mathfrak{U}$ -small sets is denoted by $(\text{Set}_{\mathfrak{U}})$. Note that by a skeleton argument, $(\text{Set}_{\mathfrak{U}})$ is equivalent to the category of sets contained as elements in $\mathfrak{U}$, denoted by $(\text{Set}_{\mathfrak{U}}^0)$.*

A small category C is said to be $\mathfrak{U}$ -small if $\text{Mor } C$ is $\mathfrak{U}$ -small. A category C is said to be essentially $\mathfrak{U}$ -small if it is equivalent to a $\mathfrak{U}$ -small category, or, equivalently, if it has a $\mathfrak{U}$ -small skeleton.

Remark 1.5 The category $(\text{Set}_{\mathfrak{U}}^0)$ is $\mathfrak{V}$ -small since $\coprod_{(x,y) \in \mathfrak{U} \times \mathfrak{U}} (\text{Set}_{\mathfrak{U}}^0)(x, y)$ is an element of $\mathfrak{V}$. Hence the category $(\text{Set}_{\mathfrak{U}})$ is essentially $\mathfrak{V}$ -small.

Given categories C and D , we denote the category of functors mapping from C to D by $\llbracket C, D \rrbracket$.

Lemma 1.6 *If C and D are $\mathfrak{U}$ -small, then $\llbracket C, D \rrbracket$ is $\mathfrak{U}$ -small. If C and D are essentially $\mathfrak{U}$ -small, then $\llbracket C, D \rrbracket$ is essentially $\mathfrak{U}$ -small.*

Using induced equivalences, it suffices to prove the first assertion. But then

$$\text{Ob } \llbracket C, D \rrbracket \subseteq \mathfrak{P}(\text{Mor } C \times \text{Mor } D)$$

is $\mathfrak{U}$ -small. Moreover, given $f, g \in \llbracket C, D \rrbracket$, the set

$$\llbracket C, D \rrbracket(f, g) \subseteq \coprod_{x \in \text{Ob } C} D(fx, gx)$$

is $\mathfrak{U}$ -small. Hence $\text{Mor } \llbracket C, D \rrbracket$ is $\mathfrak{U}$ -small.

Lemma 1.7 *Given an essentially $\mathfrak{U}$ -small category C , the category*

$$C^{\wedge \mathfrak{U}} := \llbracket C^o, (\text{Set}_{\mathfrak{U}}) \rrbracket$$

of $(\text{Set}_{\mathfrak{U}})$ -valued presheaves over C is essentially $\mathfrak{V}$ -small. Sometimes we write $C^{\wedge} = C^{\wedge \mathfrak{U}}$ if the universe is unambiguous. Likewise, the category

$$C^{\vee \mathfrak{U}} := \llbracket C, (\text{Set}_{\mathfrak{U}}) \rrbracket$$

of $(\text{Set}_{\mathfrak{U}})$ -valued copresheaves over C is essentially $\mathfrak{V}$ -small. Sometimes we write $C^{\vee} = C^{\vee \mathfrak{U}}$.

This follows from (1.5, 1.6).

Definition 1.8 *Given an essentially $\mathfrak{U}$ -small category C , we have the Yoneda embedding*

$$\begin{array}{ccc} C & \xrightarrow{y_C} & C^{\wedge \mathfrak{U}} \\ x & \longrightarrow & \mathcal{C}(-, x) \end{array}$$

and the co-Yoneda embedding

$$\begin{array}{ccc} C & \xrightarrow{y'_C} & C^{\vee \mathfrak{U}} \\ x & \longrightarrow & \mathcal{C}(x, -). \end{array}$$

For a functor $C \xrightarrow{f} D$, we denote

$$\begin{array}{ccc} C^{\wedge \mathfrak{U}} & \xleftarrow{f^{\wedge}} & D^{\wedge \mathfrak{U}} \\ (u \circ f^o \xrightarrow{\beta \circ f^o} u' \circ f^o) & \longleftarrow & (u \xrightarrow{\beta} u'). \end{array}$$

Given functors $C \xrightleftharpoons[g]{f} D$ and a natural transformation $f \xrightarrow{\alpha} g$, we denote by $f^{\wedge} \xleftarrow{\alpha^{\wedge}} g^{\wedge}$ the natural transformation that is given at $u \in D^{\wedge \mathfrak{U}}$ by the morphism $u \circ f^o \xleftarrow{u \circ \alpha^o} u \circ g^o$ in $C^{\wedge \mathfrak{U}}$, that evaluated at $c \in C^o$ in turn yields $ufc \xleftarrow{u\alpha c} ugc$.

Analogously for $C^{\vee \mathfrak{U}} \xleftarrow{f^{\vee}} D^{\vee \mathfrak{U}}$ and $f^{\vee} \xrightarrow{\alpha^{\vee}} g^{\vee}$.

2 The right Kan extension

Proposition 2.1 (the relative Yoneda Lemma) *Given essentially $\mathfrak{A}$ -small categories C and D , and a functor $C \xrightarrow{f} D$. Then the right adjoint $C^{\wedge \mathfrak{A}} \xrightarrow{\varphi} D^{\wedge \mathfrak{A}}$ of $C^{\wedge \mathfrak{A}} \xleftarrow{f^\wedge} D^{\wedge \mathfrak{A}}$ is given by*

$$\begin{array}{ccc}
 C^{\wedge \mathfrak{A} \wedge \mathfrak{B}} & \xrightarrow{f^{\wedge \wedge}} & D^{\wedge \mathfrak{A} \wedge \mathfrak{B}} \\
 \uparrow y_C^\wedge & & \downarrow y_D^\wedge \\
 & & D^{\wedge \mathfrak{B}} \\
 & & \uparrow \text{J} \\
 C^{\wedge \mathfrak{A}} & \xrightarrow{\varphi} & D^{\wedge \mathfrak{A}}
 \end{array}$$

Keeping the name of the functor after restricting the image to $D^{\wedge \mathfrak{A}}$, we write shorthand

$$f^\wedge \dashv y_D^\wedge \circ f^{\wedge \wedge} \circ y_C^\wedge.$$

The unit of this adjunction

$$1_{D^\wedge} \xrightarrow{\varepsilon} \varphi \circ f^\wedge$$

at $u \in D^{\wedge \mathfrak{A}}$, i.e.

$$u \xrightarrow{\varepsilon u} C^\wedge(f^{\wedge, o} \circ y_D^o(-), u \circ f^o),$$

applied to $d \in D$, is given by

$$\begin{aligned}
 ud & \xrightarrow{\varepsilon ud} C^\wedge(D(f^o(-), d), u \circ f^o) \\
 x & \longrightarrow (x)\varepsilon ud,
 \end{aligned}$$

where the natural transformation $(x)\varepsilon ud$ sends at $c \in C$

$$\begin{aligned}
 D(fc, d) & \xrightarrow{(x)\varepsilon ud} ufc \\
 a & \longrightarrow (a) \left[((x)\varepsilon ud)c \right] = (x)(ua^o).
 \end{aligned}$$

The counit of this adjunction

$$f^\wedge \circ \varphi \xrightarrow{\eta} 1_{C^\wedge}$$

at $v \in C^\wedge$, i.e.

$$C^\wedge(f^{\wedge, o} \circ y_D^o \circ f^o(-), v) \xrightarrow{\eta v} v,$$

applied to $c \in C$, is given by

$$\begin{aligned}
 C^\wedge(D(f^o(-), fc), v) & \xrightarrow{\eta vc} vc \\
 \xi & \longrightarrow (\xi)\eta vc = (1_{fc})\xi c
 \end{aligned}$$

Since the set

$$c^\wedge(D(f(-), d), v) \subseteq \prod_{c \in \text{Ob } C} (\text{Set}_{\mathfrak{U}})(D(fc, d), vc)$$

is $\mathfrak{U}$ -small, φ exists as the factorization of $y_D^\wedge \circ f^{\wedge\wedge} \circ y_{C^\wedge}$ over the inclusion $D^{\wedge\mathfrak{U}} \subseteq D^{\wedge\mathfrak{V}}$.

Various compatibilities need to be verified to ensure the well-definedness of ε and η ⁽¹⁾.

We have to show that $(f^\wedge \circ \varepsilon)(\eta \circ f^\wedge) = 1_{f^\wedge}$. Suppose given $u \in D^\wedge$, $c \in C$ and $x \in (f^\wedge u)c = u(fc)$. We obtain

$$\begin{aligned} (x)((f^\wedge \circ \varepsilon)uc)((\eta \circ f^\wedge)uc) &= (x)(\varepsilon u(fc))(\eta(u \circ f^o)c) \\ &= (1_{fc}) \left[((x)(\varepsilon u(fc)))c \right] \\ &= (x)(u1_{fc}^o) \\ &= x. \end{aligned}$$

We have to show that $(\varepsilon \circ \varphi)(\varphi \circ \eta) = 1_\varphi$. Suppose given $v \in C^{\wedge\mathfrak{U}}$ and $d \in D$. Note that $\varphi v = c^\wedge(f^{\wedge, o} \circ y_D^o(-), v)$ and therefore $\varphi v d = c^\wedge(D(f^o(-), d), v)$. The application $((\varepsilon \circ \varphi)vd)((\varphi \circ \eta)vd)$ writes

$$c(D(f^o(-), d), v) \xrightarrow{\varepsilon(\varphi v)d} c^\wedge(D(f^o(-), d), c^\wedge(f^{\wedge, o} \circ y_D^o \circ f^o(-), v)) \xrightarrow{(-)\eta v} c^\wedge(D(f^o(-), d), v)$$

An element $\xi \in \varphi v d$, i.e. $D(f^o(-), d) \xrightarrow{\xi} v$, is thus mapped to the composite

$$D(f^o(-), d) \xrightarrow{(\xi)\varepsilon(\varphi v)d} c^\wedge(f^{\wedge, o} \circ y_D^o \circ f^o(-), v) \xrightarrow{\eta v} v.$$

¹ Given $u \in D^{\wedge\mathfrak{U}}$, we need to see that $u \xrightarrow{\varepsilon u} \varphi \circ f^\wedge(u)$ is a natural transformation. Suppose given $d' \xrightarrow{a} d$ in D . We have to show that for any $x \in ud$

$$(x)(\varepsilon ud)((\varphi \circ f^\wedge)ua^o) = (x)(ua^o)(\varepsilon ud')$$

is an equality of natural transformations from $D(f^o(-), d')$ to $u \circ f^o$. At $c \in C$, the element $fc \xrightarrow{a'} d'$ is mapped by the left hand side to $(x)(u(a'a^o))$, and by the right hand side to $(x)(ua^o)(ua'^o)$.

We need to see that $1_{D^{\wedge\mathfrak{U}}} \xrightarrow{\varepsilon} \varphi \circ f^\wedge$ is a natural transformation. Suppose given $u \xrightarrow{s} u'$ in $D^{\wedge\mathfrak{U}}$. We have to show that for any $d \in D$ and any $x \in ud$

$$(x)(sd)(\varepsilon u'd) = (x)(\varepsilon ud)((\varphi \circ f^\wedge(s))d)$$

is an equality of natural transformations from $D(f^o(-), d)$ to $u' \circ f$. At $c \in C$, the element $fc \xrightarrow{a} d$ is mapped by the left hand side to $(x)(sd)(u'a^o)$, and by the right hand side to $(x)(ua^o)(s(fc))$.

Given $v \in C^{\wedge\mathfrak{U}}$, we need to see that $f^\wedge \circ \varphi(v) \xrightarrow{\eta v} v$ is a natural transformation. Suppose given $c' \xrightarrow{b} c$ in C . We have to show that for any $\xi \in (f^\wedge \circ \varphi(v))c = c^\wedge(D(f^o(-), fc), v)$ we have

$$(x)(\eta vc)(vb^o) = (x)((f^\wedge \circ \varphi(v))b^o)(\eta vc').$$

The left hand side yields $(1_{fc})(\xi c)(vb^o)$, the right hand side yields $(fb)\xi c'$.

We need to see that $f^\wedge \circ \varphi \xrightarrow{\eta} 1_{C^{\wedge\mathfrak{U}}}$ is a natural transformation. Suppose given $v \xrightarrow{t} v'$ in $C^{\wedge\mathfrak{U}}$. We have to show that for any $c \in C$ and any $\xi \in (f^\wedge \circ \varphi(v))c = c^\wedge(D(f^o(-), fc), v)$ we have

$$(\xi)(\eta vc)(tc) = (\xi)((f^\wedge \circ \varphi(t))c)(\eta v'c).$$

The left hand side yields $(1_{fc})(\xi c)(tc)$. The right hand side yields $(1_{fc})((\xi t)c)$.

Now suppose given $c \in C$ and $fc \xrightarrow{a} d$. We have

$$\begin{array}{llll}
D(fc, d) & \xrightarrow{((\xi)\varepsilon(\varphi v)d)c} & C^\wedge(D(f^o(-), fc), v) & \xrightarrow{\eta vc} \quad vc \\
a & \longrightarrow & (\xi)((\varphi v)a^o) & \\
& & = (\xi)_{C^\wedge}(D(f^o(-), a), v) & \\
& & = D(f^o(-), a)\xi & \longrightarrow (1_{fc})D(fc, a)(\xi c) \\
& & & = (a)(\xi c),
\end{array}$$

whence $\xi((\varepsilon \circ \varphi)vd)((\varphi \circ \eta)vd) = \xi$.

Remark 2.2 (the absolute Yoneda Lemma) *In case $f = 1_C$, we obtain $1 \dashv y_C^\wedge \circ y_{C^\wedge}$, which by uniqueness of the right adjoint yields the comparison isomorphism*

$$\begin{array}{ll}
C^\wedge(C(-, c), v) & \xrightarrow[\sim]{\eta vc} \quad vc \\
\xi & \longrightarrow (1_c)\xi c,
\end{array}$$

at $v \in C^{\wedge \mathfrak{U}}$ and $c \in C$, with inverse given by εvc .

Corollary 2.3 *If f is full, then the counit η of the adjunction $f^\wedge \dashv y_D^\wedge \circ f^{\wedge\wedge} \circ y_{C^\wedge}$ is a monomorphism. If f is full and faithful, then η is an isomorphism.*

3 The left Kan extension

For sake of comparison to (2.1), we rephrase the pertinent case of KAN's formula in our setting.

Let C be a $\mathfrak{U}$ -small category. Let $v \in C^{\wedge \mathfrak{U}}$, let $w \in C^{\vee \mathfrak{U}}$. We define the set $v \otimes_C w$ as the quotient of the disjoint union

$$v \times_C w := \coprod_{c \in C} vc \times wc$$

modulo the equivalence relation generated by the following relation $\sim_C$. The equivalence class of $(p, q) \in vc \times wc$, $c \in C$, shall be denoted by $p \otimes q$.

Given $(p, q) \in vc \times wc$, $(p', q') \in vc' \times wc'$, we say that $(p, q) \sim_C (p', q')$ if there exists a morphism $c \xrightarrow{a} c'$ such that

$$\begin{array}{ll}
(p')va^o & = p \\
(q)wa & = q'.
\end{array}$$

Thus the quotient map $v \times_C w \xrightarrow{\nu} v \otimes_C w$ has the following universal property. Given a map $v \times_C w \xrightarrow{\nu'} X$ such that for any morphism $c \xrightarrow{a} c'$, any $p' \in vc'$ and any $q \in wc$ we have

$$((p')va^o, q)\nu' = (p', (q)wa)\nu',$$

there exists a unique map $v \otimes_C w \xrightarrow{\tilde{v}'} X$ such that $\nu' = \nu \tilde{v}'$.

In particular, given morphisms $v \xrightarrow{m} v'$ and $w \xrightarrow{n} w'$, we obtain a map $m \otimes_C n$ that maps an element represented by $(p, q) \in vc \times wc$, $c \in C$, as follows.

$$\begin{array}{ccc} v \otimes_C w & \xrightarrow{m \otimes_C n} & v' \otimes_C w' \\ p \otimes q & \longrightarrow & (p)mc \otimes (q)nc \end{array}$$

Thus the tensor product defines a functor $C^{\wedge \mathfrak{U}} \times C^{\vee \mathfrak{U}} \xrightarrow{= \otimes_C -} (\text{Set}_{\mathfrak{U}})$. We denote the univalent tensor product functor by

$$\begin{array}{ccc} C^{\wedge \mathfrak{U}} & \xrightarrow{z_C} & C^{\vee \mathfrak{U} \vee \mathfrak{U}} \\ v & \longrightarrow & v \otimes_C -. \end{array}$$

Proposition 3.1 (the relative co-Yoneda Lemma, KAN [K 58, Th. 14.1]) *Given $\mathfrak{U}$ -small categories C and D , and a functor $C \xrightarrow{f} D$. The left adjoint $C^{\wedge \mathfrak{U}} \xrightarrow{\psi} D^{\wedge \mathfrak{U}}$ of $C^{\wedge \mathfrak{U}} \xleftarrow{f^\wedge} D^{\wedge \mathfrak{U}}$ is given by*

$$\begin{array}{ccc} C^{\vee \mathfrak{U} \vee \mathfrak{U}} & \xrightarrow{f^{\vee \vee}} & D^{\vee \mathfrak{U} \vee \mathfrak{U}} \\ \uparrow z_C & & \downarrow y_D'^\wedge \\ C^{\wedge \mathfrak{U}} & \xrightarrow{\psi} & (D^o)^{\vee \mathfrak{U}} = D^{\wedge \mathfrak{U}} \end{array}$$

For short,

$$\boxed{y_D'^\vee \circ f^{\vee \vee} \circ z_C \dashv f^\wedge.}$$

The unit of this adjunction

$$\boxed{1_{C^\wedge} \xrightarrow{\varepsilon} f^\wedge \circ \psi}$$

at $v \in C^\wedge$, i.e.

$$v \xrightarrow{\varepsilon v} v \otimes_C f^\vee \circ y_D' \circ f^o(-),$$

applied to $c \in C$, is given by

$$\begin{array}{ccc} vc & \xrightarrow{\varepsilon vc} & v \otimes_C {}_D(fc, f(-)) \\ x & \longrightarrow & x \otimes 1_{fc}. \end{array}$$

The counit of this adjunction

$$\boxed{\psi \circ f^\wedge \xrightarrow{\eta} 1_{D^\wedge}}$$

at $u \in D^\wedge$, i.e.

$$u \circ f^o \otimes_C f^\vee \circ y_D'(-) \xrightarrow{\eta u} u,$$

applied to $d \in D$, is given by

$$\begin{array}{ccc} u \circ f^o \otimes_C {}_D(d, f(-)) & \xrightarrow{\eta ud} & ud \\ p \otimes q & \longrightarrow & (p)uq^o, \end{array}$$

where $p \otimes q$ is represented by $(p, q) \in ufc \times {}_D(d, fc)$ for some $c \in C$.

Various compatibilities have to be verified to ensure the well-definedness of ε and η ⁽²⁾.

We have to show that $(\varepsilon \circ f^\wedge)(f^\wedge \circ \eta) = 1_{f^\wedge}$. Suppose given $u \in D^\wedge$, $c \in C$ and $x \in (f^\wedge u)c$. We obtain

$$\begin{aligned} (x)((\varepsilon \circ f^\wedge)uc)((f^\wedge \circ \eta)uc) &= (x)(\varepsilon(u \circ f^o)c)(\eta u(fc)) \\ &= (x \otimes 1_{fc})(\eta u(fc)) \\ &= (x)u1_{fc}^o. \\ &= x. \end{aligned}$$

We have to show that $(\psi \circ \varepsilon)(\eta \circ \psi) = 1_\psi$. Suppose given $v \in C^\wedge$, $d \in D$, $c \in C$, $s \in vc$, $t \in {}_D(d, fc)$, so that $s \otimes t \in \psi vd = v \otimes_C {}_D(d, f(-))$. We obtain

$$\begin{aligned} (s \otimes t)((\psi \circ \varepsilon)vd)((\eta \circ \psi)vd) &= (s \otimes t)(\varepsilon v \otimes {}_D(d, f(-)))(\eta(v \otimes_C f^\vee \circ y'_D)d) \\ &= ((s \otimes 1_{fc}) \otimes t)(\eta(v \otimes_C f^\vee \circ y'_D)d) \\ &= (s \otimes 1_{fc})(v \otimes_C {}_D(t, f(-))) \\ &= (s \otimes t). \end{aligned}$$

² Given $v \in C^\wedge$, we need to see that $v \xrightarrow{\varepsilon v} f^\wedge \circ \psi(v)$ is a natural transformation. Suppose given $c' \xrightarrow{b} c$. We have to show that for any $x \in vc$

$$(x)(\varepsilon vc)((f^\vee \circ \psi(v))b^o) = (x)(vb^o)(\varepsilon vc').$$

The left hand side yields $x \otimes fb$. The right hand side yields $x(vb^o) \otimes 1_{fc'}$.

We need to see that $1_{C^\wedge} \xrightarrow{\varepsilon} f^\wedge \circ \psi$ is a natural transformation. Suppose given $v \xrightarrow{t} v'$ in $C^\wedge$. For any $c \in C$ and any $x \in vc$ we have

$$(x)(tc)(\varepsilon v'c) = (x)(\varepsilon vc)(tc \otimes {}_D(fc, f(-))) = (x)tc \otimes 1_{fc}.$$

Given $u \in D^\wedge$ and $d \in D$, we need to see that ηud is a well-defined map. Suppose given $c, c' \in C$, $c \xrightarrow{b} c'$ in C and $p' \in ufc'$, $q \in {}_D(d, fc)$. Since

$$((p')uf^ob^o)uq^o = (p')u(q(fb))^o,$$

the universal property applies.

Given $u \in D^\wedge$, we need to see that $\psi \circ f^\wedge(u) \xrightarrow{\eta u} u$ is a natural transformation. Suppose given $d' \xrightarrow{a} d$. For all $c \in C$, $p \in ufc$ and $q \in {}_D(d, fc)$ we obtain

$$(p \otimes q)(\eta ud)(ua^o) = (p \otimes q)((\psi \circ f^\wedge(u))a^o)(\eta ud') = (p)u(aq)^o.$$

We need to see that $\psi \circ f^\wedge \xrightarrow{\eta} 1_{D^\wedge}$ is a natural transformation. Suppose given $u' \xrightarrow{s} u$ in $D^\wedge$. We have to show that for all $d \in D$, $c \in C$, $p \in ufc$ and $q \in {}_D(d, fc)$, we have

$$(p \otimes q)(\eta ud)(sd) = (p \otimes q)((\psi \circ f^\wedge(s))d)(\eta u'd).$$

The left hand side yields $(p)(uq^o)(sd)$. The right hand side yields $(p)(s(fc))(u'q^o)$.

Remark 3.2 (the absolute co-Yoneda Lemma) *In case $f = 1_C$, we obtain $y_D'^\vee \circ z_C \dashv 1_C^\wedge$, which by uniqueness of the left adjoint yields the comparison isomorphism*

$$\begin{array}{ccc} v \otimes_C c(c, -) & \xrightarrow[\sim]{\eta c} & vc \\ s \otimes t & \longrightarrow & (s)vt^o \end{array}$$

at $v \in C^{\wedge u}$ and $c \in C$, with inverse given by $\varepsilon v c$.

Corollary 3.3 *If f is full, then the unit ε of the adjunction $y_D'^\vee \circ f^{\vee\vee} \circ z_C \dashv f^\wedge$ is an epimorphism. If f is full and faithful, then ε is an isomorphism.*

4 References

BOURBAKI, N.

[SGA 4] *Univers*, appendice de l'exposé I de SGA 4, tome 1, SLN 269, 1972.

KAN, D. M.

[K 58] *Adjoint functors*, Trans. Am. Math. Soc. 87, pp. 294-329, 1958.

MAC LANE, S.

[ML 71] *Categories for the Working Mathematician*, GTM 5, 1971.

[ML 98] *The Yoneda Lemma*, Math. Japon. 47 (1), p. 156, 1998.

MITCHELL, B.

[M 65] *Theory of categories*, Academic Press, 1965.